

# Blog article on *Gloriosa superba*

***Gloriosa superba* in Chhattisgarh: A Botanical Marvel, Medicinal Powerhouse, and Conservation Imperative**



## Table of Contents

1. Introduction: Chhattisgarh's Fiery Lily – A Tapestry of Beauty, Healing, and Peril .....	3
2. Botanical Profile: Unveiling the Glory Lily's Unique Characteristics .....	3
Scientific Classification and Nomenclature .....	3
Morphological Features and Growth Habits .....	4
Natural Habitat and Distribution .....	5
3. Chhattisgarh's Connection: A Regional Deep Dive .....	5
Specific Distribution and Presence within Chhattisgarh.....	5
Local Names and Ethnobotanical Uses by Tribal Communities .....	5
Cultural Significance and Traditional Beliefs .....	6
4. The Healing and Hazardous: Medicinal Properties and Toxicity .....	7
Key Bioactive Compounds .....	7
Traditional Medicinal Applications .....	7
Modern Pharmacological Research and Therapeutic Potential.....	8
Understanding the Risks: Symptoms of Poisoning and First Aid Measures.....	8
Key Bioactive Compounds and Pharmacological Actions of <i>Gloriosa superba</i> .....	9
5. Cultivation and Commercial Value: A Growing Industry .....	11
Detailed Cultivation Practices in India .....	11
Economic Importance and Commercialization .....	13
Challenges and Opportunities in Sustainable Cultivation .....	14
6. Conservation Imperative: Protecting a Vulnerable Treasure .....	14
Global and National Conservation Status.....	14
Specific Threats in Chhattisgarh .....	15
Current Conservation Efforts and Future Strategies.....	16
Conservation Status of <i>Gloriosa superba</i> .....	16
7. Conclusion: Charting a Sustainable Future for <i>Gloriosa superba</i> .....	17
8. References .....	18

# 1. Introduction: Chhattisgarh's Fiery Lily – A Tapestry of Beauty, Healing, and Peril

*Gloriosa superba*, commonly known by a plethora of evocative names such as Gloriosa lily, glory lily, fire lily, flame lily, climbing lily, creeping lily, cat's claw, or tiger's claw, is a visually striking plant recognized globally for its vibrant appearance. Its scientific name, derived from the Latin 'gloriosus' (glorious) and 'superba' (superb), aptly captures its splendid and majestic floral display. This plant holds significant cultural importance, serving as the national flower of Zimbabwe and the state flower of Tamil Nadu, India.

The narrative of *Gloriosa superba* is characterized by a profound duality: it is both a celebrated medicinal plant and a potent toxic agent. All parts of the plant, particularly its tubers, contain high concentrations of dangerous alkaloids, primarily colchicine and gloriosine, which can prove fatal if ingested in sufficient quantities. Despite this inherent danger, the plant has a long history of use in traditional medicine, where it has been employed in carefully controlled, lower doses. In Indian medicine, it is classified as one of the 'seven upavishas' or semi-poisonous drugs, signifying its capacity to cure illnesses while simultaneously posing a lethal threat if misused. The striking contrast between its glorious appearance and symbolic status on one hand, and its severe toxicity on the other, creates a compelling narrative. This inherent contradiction highlights the deep, empirical understanding of dosage and preparation that has been developed and passed down through generations in traditional medicinal systems.

This report focuses on Chhattisgarh, India, as a critical geographical context for *Gloriosa superba*. The plant is widely distributed across various tribal regions within the state, including Rajnandgaon, South Surguja, Bilaspur, Korba, Raigarh, Raipur, Bastar, and Dantewada districts. The plant's reported endangered status within Chhattisgarh underscores the urgent need for a detailed examination of its presence, utilization, and conservation challenges unique to this region. Chhattisgarh serves as a representative case study for understanding the intricate interplay of biodiversity, traditional knowledge, economic pressures, and conservation imperatives across India. The localized threats and conservation efforts observed here offer valuable insights into broader national and global challenges faced by high-value medicinal plants.

## 2. Botanical Profile: Unveiling the Glory Lily's Unique Characteristics

### Scientific Classification and Nomenclature

*Gloriosa superba* L. is scientifically classified under the family Colchicaceae, though it was previously categorized within Liliaceae. Its widespread recognition across India is

evident in its diverse common and vernacular names. Beyond its English appellations like Malabar Glory Lily, Tiger Claw Lily, and Glorious Lily, it is known by names such as Karihari, Langoli, and Kaliari in Hindi; Agnishikha in Kannada; Kalappaikizhangu in Tamil; Nageeti gadda in Telugu; and Langali in Sanskrit. This extensive nomenclature reflects the plant's deep integration into various linguistic and cultural landscapes, indicating centuries of close human interaction and observation. The sheer volume and diversity of these names, often describing its physical characteristics or perceived properties, underscore a nuanced, localized understanding that has been passed down through generations.

## Morphological Features and Growth Habits

*Gloriosa superba* is a highly variable, deciduous, summer-growing tuberous plant. It emerges from a fleshy, red-brown, elongated, often forked tuberous rhizome in spring. Its slender stems typically grow between 1 to 6 feet long, though they can extend up to 15 feet under optimal conditions. A distinctive feature is its climbing habit, facilitated by numerous glossy emerald green, oval- to spear-shaped leaves that taper into coiling, clinging tendrils at their tips. These tendrils allow the scrambling plant to ascend other plants or structures. The stems typically die back in late summer or fall, with the plant regenerating from its resilient tubers.

The plant produces showy, solitary flowers from mid-summer to fall, borne on long stalks in the leaf axils. Each flower features six widely separated, reflexed tepals (petals and sepals that appear similar) and six long, outward-spreading stamens. The vibrant color palette typically ranges from bright red to orange, often with yellowish bases, but can also include cream, yellow, purple-red, or bicolor variations, with colors often deepening as the flowers age. Following pollination, the flowers develop into large, fleshy, oblong fruits, which are three-valved capsules containing approximately 20 rounded red seeds.

The morphological adaptations of *Gloriosa superba*, such as its tendrils for climbing and tubers for storage and regeneration, contribute to its ecological resilience and its ability to colonize diverse habitats, including nutrient-poor soils. This inherent robustness explains its classification as a "noxious weed" in some regions. However, a critical vulnerability lies in its strong apical dominance; injury to the apical shoot meristem can result in the death of the entire plant, as dormant lateral buds do not activate to compensate for the damage. This specific fragility means that while the plant is generally hardy, it is acutely susceptible to certain forms of physical damage, particularly those inflicted by human activities like unskilled harvesting or by livestock such as cattle grazing, which significantly impacts its growth and reproductive capacity, thereby contributing to its endangered status despite its apparent hardiness.

## Natural Habitat and Distribution

*Gloriosa superba* exhibits remarkable adaptability, growing naturally in a wide variety of habitats, from forests and thickets to grasslands and even nutrient-poor soils like sand dunes. It thrives in warm, humid, and tropical conditions, preferring seasonal monsoon climates with a pronounced dry season. The plant is not resistant to frost. Its global distribution spans tropical and southern Africa, and temperate and tropical Asia, extending from China to India. Within India, it is found throughout tropical regions, from the North West Himalaya to Assam and the Deccan peninsula. Specific Indian states where it is distributed include Andhra Pradesh, Karnataka, Kerala, Maharashtra, Odisha, and Tamil Nadu.

## 3. Chhattisgarh's Connection: A Regional Deep Dive

### Specific Distribution and Presence within Chhattisgarh

*Gloriosa superba* holds a significant presence in Chhattisgarh, where it is widely distributed across the state's tribal regions. Noteworthy districts where the plant is found include Rajnandgaon, South Surguja, Bilaspur, Korba, Raigarh, Raipur, Bastar, and Dantewada. Chhattisgarh is considered a prime example of the scrub forest regions where

*G. superba* commonly occurs in India. Despite its widespread distribution, the Chhattisgarh State Medicinal Plant Board has classified

*Gloriosa superba* as an endangered species within the state, highlighting the localized urgency of its conservation. The convergence of

*G. superba* populations and diverse tribal communities in Chhattisgarh positions the state not merely as a geographical area, but as a vital repository of invaluable ethnobotanical knowledge.

### Local Names and Ethnobotanical Uses by Tribal Communities

In Chhattisgarh, *Gloriosa superba* is locally known as Kalihari in Hindi. Tribal communities in the state have historically utilized this plant for various traditional medicinal purposes. For instance, they prepare a remedy for treating cancerous wounds by boiling the plant's bulb with mustard oil, also incorporating its leaves and seeds in the process. In Raigarh, Chhattisgarh, Kalihari rhizome paste is traditionally rubbed externally on the body for three days as a treatment for leprosy. These specific, documented uses for complex conditions suggest a sophisticated, empirical understanding of the plant's potent properties, including methods of preparation and dosage that mitigate its inherent toxicity. This demonstrates the critical importance of documenting and validating such traditional knowledge, as it can offer novel leads for

modern pharmaceutical research and sustainable resource management, emphasizing the need for collaborative efforts between scientific researchers and indigenous communities.

These local applications are part of a broader pattern of *Gloriosa superba*'s utilization in Indian traditional medicine. The plant is used in 12 Indian states, including Chhattisgarh, by 36 different tribes such as the Santal, Munda, Oraon, Baiga, Gond, and Bhil communities. Its traditional applications span a wide array of 29 disease conditions, including leprosy, lice, wounds, rheumatism, snakebite, scorpion bite, gout, abdominal pain, and as an abortifacient.

## Cultural Significance and Traditional Beliefs

While *Gloriosa superba* is deeply revered in Indian culture and is often associated with the goddess Saraswati, the available research does not provide specific details on its cultural significance or rituals among tribal communities in Chhattisgarh directly related to

*Gloriosa superba*. Although various general tribal rituals and deities in Chhattisgarh are documented, such as the Sarhul festival, Bheema dev, Hareli, Holika dehan, and Bastar Dussehra, a direct link between these practices and

*Gloriosa superba* is not established in the provided material. This indicates a potential gap in documented research concerning the specific cultural integration of this plant within Chhattisgarh's tribal communities, suggesting an area ripe for future ethnobotanical inquiry.

The classification of *Gloriosa superba* as "widely distributed" in Chhattisgarh's tribal regions alongside its reported "endangered" status in the state presents an apparent contradiction. This suggests that "wide distribution" may refer to its geographical spread rather than the density or health of its populations. The endangered status, despite its presence across many districts, indicates an overall decline in population, likely driven by unsustainable harvesting due to commercial demand and habitat degradation, including cattle grazing. This situation highlights the need for targeted, localized conservation strategies that go beyond general policies. Conservation efforts in Chhattisgarh must focus on specific areas experiencing high traditional use and commercial harvesting pressure, potentially involving community-based monitoring, sustainable harvesting protocols, and local cultivation initiatives to alleviate pressure on wild populations.

## 4. The Healing and Hazardous: Medicinal Properties and Toxicity

### Key Bioactive Compounds

The potent effects of *Gloriosa superba* are primarily attributed to its highly active and toxic alkaloids, notably **colchicine** and **floriosine**, which are present in all parts of the plant but are particularly concentrated in the tubers. Other identified compounds include lumicolchicine, 3-desmethyl colchicine, beta-lumicolchicine, chelidonic acid, and salicylic acid. Beyond these, the plant also contains a range of beneficial phytochemicals such as saponin glycosides, flavonoids, tannins, steroids, polyphenols, carotenoids, and ascorbic acid, which contribute to its diverse pharmacological profile. It is notable that the colchicine yield is significantly higher in the seeds (0.7% - 0.9%) compared to the rhizomes (0.15% - 0.3%).

### Traditional Medicinal Applications

Despite its inherent toxicity, *Gloriosa superba* has been extensively utilized in traditional medicine systems across tropical Africa and Asia, including Ayurveda and various folk medicine practices, typically in lower, carefully controlled doses. Its traditional applications are wide-ranging:

- **Tubers:** Historically used as an antidote for snake bites, and for treating conditions such as gout, rheumatism, bruises, sprains, colic, chronic ulcers, haemorrhoids, cancer, impotence, nocturnal seminal emission, and leprosy. They have also been traditionally employed to induce labor pains and abortions.
- **Roots:** Considered acrid, anthelmintic, antipyretic, bitter, digestive, and expectorant, with properties that promote the expulsion of the placenta. Root paste is reported to be effective against paralysis, rheumatism, snake bites, and insect bites.
- **Leaves:** Used to treat piles, ulcers, and to aid in placenta expulsion. Historically, the juice of the leaves was used to kill head lice and as an ingredient in arrow poisons.
- **General Uses:** The plant has been applied for treating wounds, various skin-related problems, fever, inflammation, blood disorders, uterine contractions, and as a general body tonic.

Its classification as one of the 'seven upavishas' (semi-poisonous drugs) in Indian medicine underscores the critical importance of expert knowledge in its preparation and dosage, as the same compounds responsible for its therapeutic benefits can be lethally toxic if not handled with extreme precision. This highlights the concept of a "narrow

therapeutic window," where the difference between an effective dose and a toxic dose is minimal. The traditional classification reflects centuries of empirical knowledge about this critical dosage sensitivity, implying that any medicinal application, whether traditional or modern, demands rigorous precision and expertise. This also suggests that traditional preparation methods, which may involve detoxification processes, warrant further ethnopharmacological investigation.

## Modern Pharmacological Research and Therapeutic Potential

Contemporary scientific investigations have begun to validate and expand upon the traditional uses of *Gloriosa superba*. Colchicine and gloriosine are recognized as key phytochemicals for treating gout and rheumatism. Modern research has revealed several pharmacological activities:

- **Anti-inflammatory and Analgesic:** Alkaloids like colchicine demonstrate potent anti-inflammatory effects by inhibiting inflammatory mediators. Alcoholic, hydroalcoholic, and aqueous extracts from the tubers have shown significant anti-inflammatory activity.
- **Anti-cancer:** Methanolic extracts exhibit anti-oxidant and cytotoxic effects on Hep-G2 (human liver cancer) cells. Colchicine is utilized in cancer treatment due to its ability to inhibit cellular division.
- **Anti-diabetic:** Phytochemical screening indicates the presence of compounds such as saponin glycosides, flavonoids, alkaloids, tannins, and steroids, which are associated with hypoglycemic action.
- **Other Activities:** The plant exhibits anti-anxiety, anti-oxidant, antibacterial, antimicrobial, and anthelmintic properties. Colchicine is also used in clinical practices for acute pericarditis, cirrhosis of the liver, and severe constipation.

## Understanding the Risks: Symptoms of Poisoning and First Aid

### Measures

The severe toxicity of *Gloriosa superba* cannot be overstated; all parts of the plant, particularly the tubers, are extremely poisonous. Ingestion can lead to life-threatening conditions.

### Symptoms of Poisoning:

Initial symptoms typically manifest within two to six hours after ingestion. These include numbness and tingling around the mouth, a burning sensation and rawness in the throat, nausea, intense vomiting, abdominal pain, and bloody diarrhoea, often leading to severe dehydration. More critical complications can follow, such as respiratory depression, dyspnoea, shock, hypotension, marked leucopenia, thrombocytopenia, coagulation

disorders, oliguria, haematuria, confusion, seizures, and coma. Delayed manifestations, appearing one to two weeks after poisoning, include alopecia (hair loss) and dermatitis.

### First Aid and Management Principles:

Immediate medical attention is crucial in cases of suspected *Gloriosa superba* poisoning. If the patient is conscious and alert, vomiting should be induced (e.g., by tickling the back of the throat or administering syrup of ipecac, followed by water). If this is ineffective, a stomach wash-out should be performed. The patient must be admitted to a hospital immediately, and any remaining plant material or vomit should be collected for analysis. Clinical management involves careful monitoring of respiration, ensuring an adequate airway, immediate gastric lavage, treating hypotension with intravenous fluids and vasopressors, and potentially blood transfusions to support circulation. Correcting dehydration and electrolyte imbalance, monitoring renal function, and performing initial forced diuresis to enhance colchicine elimination are also vital.

The significant economic value of *Gloriosa superba* due to its medicinal compounds, coupled with its widespread use in traditional medicine, creates a substantial public health challenge. This is particularly true in regions where the plant grows wild or is cultivated without strict regulatory oversight. The economic incentive for harvesting, combined with traditional beliefs about its healing powers, can inadvertently lead to misuse or accidental poisoning if proper knowledge of identification, preparation, and dosage is lacking. This situation underscores the urgent need for public awareness campaigns, especially in rural and tribal communities, to educate individuals about the dangers of self-medication and the critical importance of seeking expert medical guidance. It also highlights the necessity of robust regulatory frameworks for harvesting, processing, and commercialization to ensure product safety and prevent fatal incidents.

## Key Bioactive Compounds and Pharmacological Actions of *Gloriosa superba*

Compound Name	Primary Source/Part	Traditional Uses	Modern Pharmacological Actions (Validated)	Key Toxic Effects
Colchicine	Tubers, Seeds, Whole Plant	Gout, Rheumatism, Snakebite, Bruises,	Anti-inflammatory, Anti-cancer (cytotoxic),	Cellular division inhibition, Diarrhoea,

Compound Name	Primary Source/Part	Traditional Uses	Modern Pharmacological Actions (Validated)	Key Toxic Effects
		Sprains, Chronic Ulcers, Haemorrhoids, Cancer, Leprosy, Inducing Labor/Abortion, Fever, Inflammation, Skin problems	Anti-diabetic, Anti-oxidant, Antimicrobial, Microtubule polymerization inhibition	Bone marrow depression, Alopecia, Nausea, Vomiting, Abdominal pain, Bloody diarrhoea, Respiratory depression, Shock, Seizures, Coma
Gloriosine	Tubers, Whole Plant	Gout, Rheumatism	Anti-inflammatory	Similar to Colchicine, contributes to overall toxicity
Lumicolchicine	Whole Plant	-	-	Toxic
Flavonoids	Tubers, Seeds	-	Anti-diabetic, Anti-oxidant	-

Compound Name	Primary Source/Part	Traditional Uses	Modern Pharmacological Actions (Validated)	Key Toxic Effects
Saponins	Tubers, Seeds	-	Anti-diabetic, Anti-anxiety	-
Tannins	Tubers, Seeds	-	Anti-diabetic	-
Steroids	Tubers, Seeds	-	Anti-diabetic	-
Polyphenols	Tubers, Seeds	-	Anti-oxidant	-
Carotenoids	Tubers, Seeds	-	Anti-oxidant	-
Ascorbic Acid	Tubers, Seeds	-	Anti-oxidant	-

## 5. Cultivation and Commercial Value: A Growing Industry

### Detailed Cultivation Practices in India

The cultivation of *Gloriosa superba* in India requires specific environmental conditions and meticulous practices to ensure successful growth and yield.

#### Climate and Soil:

The plant thrives in warm, humid, tropical regions with a distinct dry season and is intolerant to frost. Optimal soil conditions include Black Soil or Red Sandy Loam, with a pH ranging from 5.5 to 7.0, and critically, excellent drainage to prevent waterlogging.

## Land Preparation:

Thorough ploughing and harrowing are essential to achieve a fine soil tilth. Proper leveling and the establishment of an effective drainage system are crucial. Approximately 10 tonnes of Farm Yard Manure (FYM) per acre should be applied during land preparation, and beds should be prepared into ridges and furrows.

## Propagation:

While seeds can be used, plants are best propagated from 'V' shaped tubers. Seed propagation is a slow process, taking 3-4 years to bloom, and is challenging due to hard seed coats and a dormancy period of 6-9 months. Tubers are brittle and require careful handling during planting. The optimal tuber weight for planting is 50-60g, as smaller rhizomes may not flower in the first year.

## Planting:

The ideal planting season in India is June-July, coinciding with the Kharif season. Tubers should be planted horizontally 2-4 inches deep in furrows 6-8 cm deep, with a plant-to-plant spacing of 20-30 cm, or a general spacing of 60 x 45 cm. Approximately 16,000 tubers per acre are required, accounting for a 7% mortality rate, or about 600 kg of rhizomes for 12,000-13,000 plants per acre. Closer spacing can promote cross-pollination and improve fruit set.

## Support System:

As a climbing plant, *Gloriosa superba* necessitates a support system. A thin-wire trellis or allowing it to scramble through other plants can be effective. Farmers commonly use 4-foot wooden sticks planted 15 feet apart, with 3-4 parallel steel wires and sorghum straw to facilitate climbing. It is important to handle stems carefully, as excessive manipulation can cause damage and inhibit growth.

## Manure and Fertilization:

For medicinal plants like *Gloriosa superba*, organic manures and fertilizers are recommended, with a strict avoidance of chemical fertilizers and pesticides. Vermicompost, Neem Cake (as an organic insecticide), Gypsum (a soil conditioner), and Trichoderma (a fungicide) are suggested. An application of 10 tonnes of compost per acre is also recommended. A fertilizer dose of 60 kg N, 25 kg P<sub>2</sub>O<sub>5</sub>, and 40 kg K<sub>2</sub>O per hectare is advised, with P<sub>2</sub>O<sub>5</sub> and K<sub>2</sub>O applied as basal doses and nitrogen split into basal and subsequent applications.

## Irrigation:

*Gloriosa superba* is primarily a rain-fed crop, but periodic irrigation significantly enhances growth. During the initial seedling stages, frequent irrigation (every 4-7 days) is required to keep the soil soft for sprouting. Subsequently, irrigation can be reduced to 15-

day intervals. No irrigation is necessary during harvesting or after flowering. Overwatering must be avoided as it can lead to premature fruit falling.

### Weeding and Pest Management:

Frequent manual weeding and hoeing are crucial, especially during the initial growth stages. Chemical weeding should not be employed. Extreme care must be taken during weeding to avoid damaging the growing tip of the rhizomes, as such injury can prevent further sprouting. Continuous cloudy weather can favor the multiplication of

*Curvularia lunata*, which causes leaf blight, and soilborne pathogens like

*Macrophomina phaseolina* can cause dry root rot. Bio-pesticides derived from Neem, Chitrakmool, Dhatura, or Cow's urine are suggested for pest and disease management.

### Harvesting and Post-Harvesting:

Plants typically begin harvesting approximately 170-180 days after planting. Capsules are ready for harvest when they transition from light green to dark green, appear shrunken, become lighter in weight, and produce a cracking sound when pressed. Tubers are harvested after 5-6 years of transplanting. Post-harvest, tubers, seeds, and pericarp are cleaned, washed, and air-dried in the shade before being packed in airtight bags to minimize spoilage and extend shelf-life.

### Economic Importance and Commercialization

*Gloriosa superba* is recognized as a high-value medicinal plant and an expensive export commodity. The National Medicinal Plant Board of India has included it in its list of 32 nationally prioritized medicinal plants, largely due to the commercially important alkaloids, colchicine and colchicoside, it contains. Notably,

*G. superba* yields higher quantities of colchicine than *Colchicum*, making it a preferred source for this compound. The plant offers significant economic returns, with an average net return per acre reported at approximately Rs. 1,499,002 in 2019, classifying it as a cash crop comparable to sugarcane and cotton. Commercialization of

*G. superba* began with tuber exports, followed by seed exports starting in 1984.

The significant economic value of *G. superba* creates a powerful, market-driven incentive for harvesting, which, when unsustainable, directly contributes to its critically vulnerable status. This highlights a fundamental conflict: the very properties that make the plant valuable also drive its depletion in the wild. However, this economic value can also be strategically leveraged for conservation. By promoting and supporting commercial cultivation, the economic incentive can be redirected from wild harvesting to controlled farming. This suggests that sustainable economic models are not merely an add-on but a crucial component of effective conservation strategies, offering a pathway to meet demand without further depleting wild populations.

## Challenges and Opportunities in Sustainable Cultivation

Despite its economic allure, the commercialization of *Gloriosa superba* faces critical challenges. Continuous and unsustainable collection of tubers from natural habitats for commercial exploitation has led to its scarcity in Indian forests. This pressure is reflected in its critically vulnerable status nationally and its endangered status in Chhattisgarh. A unique biological vulnerability of the plant is its high sensitivity to damage due to strong apical dominance; injury to the apical shoot meristem results in the death of the entire plant, as dormant lateral buds do not activate to compensate. This specific trait means that even minor damage to the primary growth point can be fatal for the individual plant. Mass destruction has been observed due to cattle grazing in natural habitats, a significant threat exacerbated by this biological characteristic. Furthermore, conventional propagation methods are limited by poor seed germination (due to hard seed coats and dormancy) and slow tuber multiplication (each plant produces a maximum of two daughter tubers), restricting its ability to replenish wild populations. These combined factors, coupled with the immense pressure from national and international drug markets, necessitate a concerted shift towards sustainable cultivation practices to meet demand and alleviate pressure on wild populations. This implies that protective measures, such as fencing, controlled grazing, and educating local communities on careful handling, are as critical as reducing overall collection pressure.

## 6. Conservation Imperative: Protecting a Vulnerable Treasure

### Global and National Conservation Status

*Gloriosa superba*'s threatened status is formally recognized at multiple levels. It is listed as an endangered species by the International Union for Conservation of Nature (IUCN), placing it in the "Red Data Book". The IUCN classification system categorizes species based on their extinction risk, ranging from Extinct to Least Concern. Within India, the Ministry of Environment, Government of India, has classified

*Gloriosa superba* as "critically vulnerable". Further emphasizing its national importance and the urgency of its conservation, the National Medicinal Plant Board of India has included it in its list of 32 nationally prioritized medicinal plants.

This significant discrepancy, where *Gloriosa superba* is globally listed as "Least Concern (LC)" by IUCN, but "critically vulnerable" nationally in India and "endangered" in Chhattisgarh, indicates that the global status may not accurately reflect the severe localized threats. While the species might be widespread globally, intense commercial exploitation and specific regional pressures, such as cattle grazing in Chhattisgarh, are causing rapid declines in key biodiversity hotspots. This situation highlights that

conservation efforts need to be highly localized and context-specific, as broader assessments can mask urgent regional crises. It also suggests potential data gaps in global assessments regarding the true extent of regional population declines and threats.

## Specific Threats in Chhattisgarh

Within Chhattisgarh, *Gloriosa superba* faces acute and localized threats, contributing to its endangered status as reported by the Chhattisgarh State Medicinal Plant Board.

### Overexploitation:

Continuous and unsustainable collection of tubers and seeds from natural habitats for commercial exploitation, driven by the immense pressure from national and international drug markets, has led to its scarcity. This mirrors a broader trend in India, where over 90% of medicinal plants face threats from excessive and unskilled harvesting.

### Habitat Degradation:

While not explicitly detailed for Chhattisgarh in the provided material, general habitat loss, both natural and human-induced, is a common threat to red-listed species globally.

### Cattle Grazing:

A particularly significant threat identified in Chhattisgarh's Rajnandgaon district is the mass destruction of *G. superba* due to cattle grazing. This problem is exacerbated by the plant's high sensitivity to damage at its apical meristem, which, due to its strong apical dominance, leads to the death of the entire plant when injured.

### Poor Natural Propagation:

The plant's inherently slow natural multiplication contributes significantly to its vulnerability. This is characterized by poor seed germination rates (due to hard seed coats and dormancy for 6-9 months) and slow tuber dormancy, with each plant producing a maximum of two daughter tubers annually. This limited reproductive capacity makes wild populations highly susceptible to over-collection and slow to recover.

The endangerment of *G. superba* is not a simple problem; it represents a complex interplay of inherent biological traits, powerful market forces, and potentially inadequate regulatory or conservation frameworks. The plant's slow recovery rate means that even moderate harvesting can have significant long-term impacts. The high economic value creates a strong incentive for illegal or unsustainable harvesting, which can overwhelm existing conservation efforts. This situation underscores that effective conservation requires a multi-faceted approach: addressing biological limitations through propagation

research, managing economic demand through sustainable cultivation and fair trade, and strengthening policy and enforcement to protect wild populations, especially in vulnerable regions like Chhattisgarh.

## Current Conservation Efforts and Future Strategies

Effective conservation of *Gloriosa superba* necessitates a multi-pronged approach, encompassing both **in-situ** (on-site, within natural habitats) and **ex-situ** (off-site, in botanical gardens, seed banks, or controlled cultivation) methods.

### Biotechnological Approaches:

Advanced techniques play a crucial role in conservation and mass multiplication. Micropropagation, callus induction, and the use of apical buds and nodal segments have been successfully employed by researchers. These in-vitro methods are vital for overcoming the limitations of conventional propagation, such as slow growth from tubers and seeds, and can alleviate pressure on wild populations, thereby preserving genetic diversity.

### Sustainable Cultivation:

Identifying and promoting the cultivation of elite germplasms with high colchicine content is paramount. This strategy supports commercial cultivation, ensuring a continuous supply for the herbal medicine industry and reducing reliance on wild harvesting.

### Collaborative Efforts:

Bridging ancient wisdom with modern science through collaborative efforts among traditional healers, botanists, and pharmacologists is essential. Such partnerships ensure that conservation strategies are culturally sensitive, scientifically informed, and ultimately more effective.

## Conservation Status of *Gloriosa superba*

Level of Assessment	Conservation Status	Source/Authority	Primary Contributing Threats
Global	Least Concern (LC)	IUCN Red List	-

Level of Assessment	Conservation Status	Source/Authority	Primary Contributing Threats
National (India)	Critically Vulnerable	Ministry of Environment, Govt. of India	Overexploitation/Over-collection (for colchicine), Habitat Loss/Degradation, Poor Natural Propagation (slow seed germination, tuber dormancy)
State (Chhattisgarh)	Endangered	Chhattisgarh State Medicinal Plant Board	Overexploitation/Over-collection (for colchicine), Cattle Grazing, Poor Natural Propagation (slow seed germination, tuber dormancy)

## 7. Conclusion: Charting a Sustainable Future for *Gloriosa superba*

*Gloriosa superba* stands as a remarkable botanical entity, celebrated for its striking aesthetic appeal, its profound traditional and modern medicinal value, and its significant economic role as a high-value medicinal plant. However, this multifaceted importance is inextricably linked to its inherent toxicity and, critically, its vulnerable status, particularly within regions like Chhattisgarh.

The future of *Gloriosa superba* hinges on a delicate balance between its utilization for economic and medicinal benefits and its urgent conservation. Uncontrolled exploitation poses a severe threat to its survival, especially in regions such as Chhattisgarh, where it is currently endangered. The plant's survival is deeply intertwined with human livelihoods, traditional practices, and global market dynamics. This implies that effective conservation solutions must extend beyond purely ecological interventions; they must also address socio-economic factors, promote equitable benefit-sharing from commercialization, respect indigenous rights and knowledge, and foster a sense of shared responsibility.

The path forward necessitates concerted and collaborative action. Integrated approaches involving rigorous scientific research, particularly in sustainable cultivation techniques and pharmacological validation, are paramount. This must be coupled with the implementation of responsible and sustainable harvesting practices and meaningful community engagement, especially with tribal communities who possess invaluable traditional knowledge regarding the plant. Given its prominence and the well-documented challenges it faces, *Gloriosa superba* can serve as a flagship species for raising awareness and galvanizing action for the conservation of other threatened medicinal plants in India. Its story encapsulates many of the broader issues confronting medicinal plant biodiversity, demonstrating that sustainable utilization and robust conservation can indeed coexist. Such concerted efforts are essential for ensuring the long-term survival of *Gloriosa superba* and for the continued benefit of both people and nature in Chhattisgarh and beyond.

## 8. References

1. Ade, R., & Rai, V. R. (2012). *Gloriosa superba* L. : An Endangered Medicinal Plant. CABI Digital Library. <https://www.cabidigitallibrary.org/doi/pdf/10.5555/20163025396>
2. Ashokkumar, K. (2015). *Gloriosa superba* (L.): A Brief Review of its Phytochemical Properties and Pharmacology. ResearchGate. [https://www.researchgate.net/publication/287294241\\_Gloriosa\\_superba\\_L\\_A\\_Brief\\_Review\\_of\\_its\\_Phytochemical\\_Properties\\_and\\_Pharmacology](https://www.researchgate.net/publication/287294241_Gloriosa_superba_L_A_Brief_Review_of_its_Phytochemical_Properties_and_Pharmacology)
3. Dhanabalan, S. P., Johnson, I., Kumaresan, P. V., Kandasamy, R., Natesan, S., Periyannan, S., & Muthusamy, K. (2024). *Characterization and Pathogenicity of Soilborne Pathogens in Gloriosa superba: Effects of Single- and Multiple-Pathogen Coinfection on Disease Responses*. *Plant Disease*, 108(11). <https://apsjournals.apsnet.org/doi/abs/10.1094/PDIS-03-24-0496-RE>
4. eFloraofIndia. (n.d.). *Gloriosa superba*. Retrieved from <https://efloraofindia.com/efi/gloriosa-superba/>
5. India Flora. (n.d.). *Gloriosa superba* L. Retrieved from <https://indiafloras.iisc.ac.in/FloraPeninsular/plants.php?name=Gloriosa%20superba>
6. Jain, R., & Jain, S. K. (2010). *Traditional medicinal plants as anticancer agents from Chhattisgarh, India*. Ayurpharm. <https://www.ayurpharm.com/imagesuser/ayurpharm132.pdf>
7. Jana, S., & Shekhawat, G. S. (2011). *Critical review on medicinally potent plant species: Gloriosa superba*. ResearchGate.

<https://www.researchgate.net/publication/47717335> Critical review on medicinally potent plant species *Gloriosa superba*

8. Kandasamy, R., & Johnson, I. (2021). *Nature's Healing Emissary – A Deep Dive into Medicinal Marvels*. BITSathy. <https://www.bitsathy.ac.in/natures-healing-emissary-a-deep-dive-into-medicinal-marvels/>
9. Maroyi, A., & van der Maesen, J. (2013). *Gloriosa superba (Colchicaceae): ethnobotany and economic importance*. CABI Digital Library. <https://www.cabidigitallibrary.org/doi/abs/10.5555/20133318893>
10. Missouri Botanical Garden. (n.d.). *Gloriosa superba 'Rothschildiana'*. Retrieved from <https://www.missouribotanicalgarden.org/PlantFinder/PlantFinderDetails.aspx?taxonid=245907&isprofile=0&>
11. Nagendra Kumar Chandrawanshi, K. L. Tiwari, S. K. Jadhav, & Afaque Quraishi. (2013). *Threats Facing by The Critically Vulnerable Medicinal Plant *Gloriosa superba* L. At Natural Habitat Major Insect Pest and Human Interference*. ResearchGate. <https://www.researchgate.net/publication/290123275> Threats Facing by The Critically Vulnerable Medicinal Plant *Gloriosa superba* L At Natural Habitat Major Insect Pest and Human Interference
12. Pareta, S. K., & Pareta, S. K. (2015). *Tribals or communities associated with this grove have deep faith on their deity*. All Subject Journal. <https://www.allsubjectjournal.com/assets/archives/2015/vol2issue2/135.3.pdf>
13. Pawar, P., Kadam, V., & Jadhav, A. (2021). *Gloriosa superba Linn.: A Review*. World Journal of Pharmaceutical Research, 10(12), 220-230. <https://www.wjpmr.com/download/article/91122021/1640932346.pdf>
14. Robsplants. (n.d.). *Gloriosa superba*. Retrieved from <http://www.robsplants.com/plants/GloriSuper>
15. Samantara, K., Dash, U., & Dash, D. (2014). *Gloriosa Superba - an Endangered plant spotted for the first time from forest of Tpchanchi Hazaribag, Jharkhand, India*. ResearchGate. <https://www.researchgate.net/publication/267861499> *Gloriosa Superba - an Endangered plant spotted for the first time from forest of Tpchanchi Hazaribag Jharkhand India*
16. The Marginpar. (n.d.). *Enchantingly beautiful, *Gloriosa**. Retrieved from <https://www.marginpar.com/stories/enchantingly-beautiful-gloriosa>

17. The International Programme on Chemical Safety (IPCS). (n.d.). *PIM 245 Gloriosa superba*. Inchem. <https://www.inchem.org/documents/pims/plant/pim245.htm>
18. The University of Wisconsin-Madison, Division of Extension. (n.d.). *Gloriosa Lily (Gloriosa superba)*. Retrieved from <https://hort.extension.wisc.edu/articles/gloriosa-lily-gloriosa-superba/>
19. Tamil Nadu Agricultural University. (n.d.). *Medicinal & Aromatic Plants - Gloriosa superba*. Agritech. [https://agritech.tnau.ac.in/banking/pdf/Medicinal%20Aromatic\\_Medicinal%20&%20Aromatic%20Plants.pdf](https://agritech.tnau.ac.in/banking/pdf/Medicinal%20Aromatic_Medicinal%20&%20Aromatic%20Plants.pdf)
20. Vaishnavi, B. A., & Hemalatha, S. (2019). *A Review on Pharmacological Properties of Glory Lily (Gloriosa superba L.)*. International Journal of Current Microbiology and Applied Sciences, 8(2), 1989-1997. <https://www.ijcmas.com/8-2-2019/B.A.%20Vaishnavi,%20et%20al.pdf>
21. Verma, D., Singh, S., & Singh, A. K. (2021). *Conservation strategies for threatened medicinal plants of India: An overview*. PMC. <https://pmc.ncbi.nlm.nih.gov/articles/PMC8148398/>
22. Verma, R. K., & Singh, P. (2019). *Kalihari (Gloriosa superba Linn.)*. HCMS. <https://www.hcms.org.in/pdf/kalihari-eng.pdf>
23. Yadav, R., Kumar, A., & Singh, P. (2021). *Gloriosa superba L. (Colchicaceae): A high-value medicinal plant indigenous to Africa and Southeast Asia*. PMC. <https://pmc.ncbi.nlm.nih.gov/articles/PMC7866646/>